

THE DISEASE AND ITS EPIDEMIOLOGY

A. Agent Polio is caused by poliovirus (genus Enterovirus), which has three serotypes. Type 1 virus is most frequently involved in epidemics and is most often isolated from paralytic cases of poliomyelitis. Type 3 and, to a lesser degree, type 2 viruses can also cause paralysis. All three are contained in the vaccine. B. Clinical Description Infection with poliovirus results in a spectrum of manifestations. The overwhelming majority of infections (95%) are clinically inapparent. About 4 – 8% of infected individuals will experience non-specific viral symptoms, such as a low-grade fever, headache, sore throat, nausea, abdominal pain, constipation, diarrhea, and/or vomiting (referred to as abortive disease). About 1–5% of infections will result in aseptic meningitis, involving stiffness of the back, neck and/or legs, at times with paresthesias, a few days after the minor illness has resolved. Only about 0.1 – 1% of infections will progress to acute flaccid paralysis (AFP) with loss of reflexes in the involved limbs, usually with fever present (paralytic poliomyelitis). Please note, today in the U.S., the most common cause of AFP is Guillain-Barré syndrome (GBS). Polio has been eradicated from the western hemisphere. Progression to paralytic poliomyelitis usually occurs within 2 – 4 days and rarely continues after the fever subsides. Spinal paralysis is typically asymmetric and more severe proximally than distally. Paralysis may compromise respiration and swallowing. After the acute episode, many patients recover at least some muscle function and prognosis for recovery can usually be established within 6 months after onset of paralytic disease. Between 2% – 10% of paralytic infections are fatal, due to respiratory compromise. Risk factors for paralytic disease include larger inoculum of poliovirus, increasing age, pregnancy, strenuous exercise, tonsillectomy, and intramuscular injections administered while the patient is infected with poliovirus. C. Reservoirs Humans are the only known host. D. Modes of Transmission The principal mode of transmission is person-to-person by the fecal-oral route. Transmission via oral secretions, such as saliva, is very uncommon but may account for some cases. In rare instances, the virus may be transmitted by contaminated sewage or water. Asymptomatic individuals, especially children, comprise a significant source of infections. No reliable evidence of spread by insects exists. No long-term carrier state is known. In temperate climates, poliovirus infections are most common in the summer and fall. Report Immediately by phone

Page 2

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04 Poliomyelitis
2E. Incubation period Abortive (non-paralytic) polio: The incubation period is usually 3 - 6 days. Paralytic polio: The incubation period is usually 7 - 21 days, with a range of 3 - 35 days. F. Period of Communicability or Infectious Period The period of communicability is not precisely defined. It appears to be greatest 7 - 10 days before and after onset of clinical symptoms, when poliovirus is present in the throat and excreted in the highest quantities in the feces. Poliovirus can continue to be shed in the feces for 4 - 6 weeks. Rarely, excretion of poliovirus >6 months after infection has been found in asymptomatic, immunodeficient persons. Poliovirus can be found in throat secretions as early as 36 hours and in the feces 72 hours after exposure to infection in both symptomatic and asymptomatic cases. G. Epidemiology Prior to the widespread use of polio vaccine, poliomyelitis occurred worldwide. Polio was epidemic in the U.S.

for the first half of the 20th century with over 20,000 cases of paralytic disease in 1952. The first inactivated poliovirus vaccine (IPV) was introduced in 1955, monovalent oral poliovirus vaccine (OPV) in 1961, trivalent (TOPV) in 1963, and enhanced inactivated poliovirus vaccine (eIPV) in 1987. After the introduction of vaccination, the reported number of cases of poliomyelitis in the U.S. dropped to <100 in 1965 and <10 cases in 1973. The last cases of indigenously-transmitted wild-type poliovirus in the U.S. were in 1979. The last case of wild-type polio disease in the Western Hemisphere was detected in Peru in 1991. The Western Hemisphere was declared free from indigenous wild-type poliovirus transmission in 1994. Poliomyelitis is now on the verge of worldwide eradication. Almost all of the world is now considered polio-free. Worldwide efforts to eradicate polio in the few countries where the disease is still endemic are underway. Strategies include: (1) achieving and maintaining high vaccination coverage among infants < 1 year old; (2) developing sensitive surveillance systems for AFP including a laboratory network; (3) conducting National Immunization Days; (4) and conducting “mopping-up” campaigns to directly target geographic areas known to be high risk for polio transmission. The number of countries where poliovirus continues to be isolated has decreased substantially, with Afghanistan, Egypt, India, Niger, Nigeria, and Pakistan the major areas of wild-type virus circulation. Due to the success of global efforts towards eradication and the elimination of indigenously transmitted disease in the Western Hemisphere, cases of paralytic poliomyelitis in the industrialized countries have become almost non-existent. During the period 1980–94, there was an average of 8–9 cases of paralytic polio reported annually in the U.S. Most of these cases were vaccine-associated paralytic poliomyelitis (VAPP), which can occur after receipt of OPV. This very rare disease accounted for an average of 8 reported cases per year in the US, during the period 1980–94 (or 1 case for every 2.4 million doses of OPV distributed). The risk for VAPP is highest after receipt of the first dose of poliovirus vaccine, occurring at one case per 750,000 doses distributed. Since 1986, the only cases of paralytic poliomyelitis occurring in the US have been vaccine-associated. In January 1997, in an effort to reduce the risk of VAPP, a sequential polio vaccination schedule (IPV for doses 1 and 2, OPV for doses 3 and 4) was recommended in the US. With the continued success of worldwide efforts to eradicate poliovirus and in the interest of completely eliminating the occurrence of VAPP, an all-IPV immunization schedule was initiated on January 1, 2000 in the US. Despite the great achievement in polio eradication in the U.S., vigilance is needed because of the possibility of importation of wild poliovirus from areas of the world where it is endemic. The importation of wild poliovirus from polio-endemic regions of the world may occur among under-immunized (1) tourists, (2) immigrants revisiting their countries of origin, or (3) members of religious groups opposed to vaccination, regardless of travel history. In 1992–93 an outbreak occurred in the Netherlands among members of a religious group that refuse immunization. Poliovirus has also been isolated from members of a similar religious group in Canada, although no cases of disease occurred.

Page 3

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04 Poliomyelitis
3 In 2003, 682 cases were reported worldwide, with approximately 90% of the cases
from Nigeria, India and Pakistan. H. Bioterrorism Potential None. 2) DISEASE

REPORTING AND CASE INVESTIGATION A. Purpose of Surveillance and Reporting • To distinguish between wild-type and vaccine-associated polio and to identify susceptible people exposed to wild-type polio. • To maintain indigenous transmission rates of wild-type poliovirus at zero. • To identify cases of VAPP that might occur secondary to immunization with OPV given in another country. B. Laboratory and Healthcare Provider Reporting Requirements Iowa Administrative Code 641-1.3(139) stipulates that the laboratory and the healthcare provider must report any suspected or confirmed cases of polio immediately. The reporting number for IDPH Center for Acute Disease Epidemiology (CADE) is (800) 362-2736, if you are calling after business hours you may call the Iowa State Patrol Office at (515) 323-4360. Laboratory Testing Services Available After communicating with IDPH, contact the University of Iowa Hygienic Laboratory bacteriology department at (319) 335-4500 for further instructions. C. Local Public Health Agency Follow-up Responsibilities Polio follow-up and case investigation is undertaken by the Local Public Health Agency (LPHA) and will be coordinated if necessary with IDPH Bureau of Disease Prevention and Immunization. Initial Question to Ask Healthcare Provider and Patient In order to assess the likelihood that a suspect case is a true case prior to laboratory testing, LPHA and/or other public health staff helping in the investigation should ask about: 1) clinical information, 2) polio immunization history of case and close contacts, 3) pertinent medical history including underlying illness/immunosuppression, 4) membership in religious/social group that might refuse immunization, 5) country of origin and length of residence in US, 6) recent history of travel (where and dates), 7) whether there were any recent out-of-town visitors (from where and dates), and 8) whether occupation entails handling of specimens that might contain poliovirus (e.g., lab work). 3) CONTROLLING FURTHER SPREAD A. Minimum Period of Isolation of a Suspect or Confirmed Case Place case on enteric precautions for six weeks after onset of symptoms or until poliovirus can no longer be recovered from feces (the number of negative specimens needed will be determined by the IDPH on a case-by-case basis). B. Minimum Period of Quarantine of Contacts None. C. Protection of Contacts of a Case Implement control measures as described below while waiting for laboratory confirmation. While indigenous transmission of wild-type poliovirus in the United States (and the Western Hemisphere as a whole) has not occurred since 1991, the importation of poliovirus from polio-endemic regions may occur among under-immunized (1) tourists, (2) immigrants revisiting their countries of origin, or (3) members of religious groups who might refuse immunization, regardless of travel history. Polio-

Page 4

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04 Polio-
 4 endemic regions include Afghanistan, Egypt, India, Niger, Nigeria, and Pakistan. An IDPH epidemiologist can help assess the likelihood of exposure to wild-type polio. OPV is still being used outside of the U.S. Vaccine-associated paralytic poliomyelitis (VAPP) should also be considered as a cause of paralysis, especially if a patient has onset of paralysis after receipt of a first dose of OPV. No control measures are indicated if the case is determined to likely be VAPP. It is also possible that the case of paralysis is due to an infectious agent other than poliovirus, such as enterovirus, or due

to some other noninfectious cause, and therefore not contagious. Therefore, it is crucial that laboratory testing be initiated to determine if the causative agent of paralysis is poliovirus and to differentiate wild-type from vaccine strain poliovirus. Identify individuals or groups who may have been exposed to the case. Also, attempt to identify the route of introduction of poliovirus into the community. To identify these groups, think in terms of “zones of exposure” and consider members of the following groups: • Household members • School/ child care associates (students/attendees and staff) • Staff and patients at medical facility where patient was cared for, especially if there was the potential for direct contact with feces or oral secretions • Religious/social groups • Sports teams and other extracurricular groups • Bus mates • Close friends • Travelers from polio-endemic regions such as Afghanistan, Egypt, India, Niger, Nigeria, and Pakistan • Any other persons who may have come in direct contact with the case’s feces or oral secretions Identify high-risk susceptibles who had contact with the case during infectious period: • Pregnant women should be referred to their obstetricians. (In child care or school settings remember to determine whether teachers, student-teachers, staff or students are pregnant.) • Immunocompromised individuals should be referred to their healthcare providers. • Infants < 6 weeks old (who are too young to have been vaccinated) should be referred to their pediatricians. Identify and vaccinate all other susceptibles > 6 weeks old with IPV (if not contraindicated). These are individuals without proof of immunity, including those with medical or religious exemptions to immunization. Proof of immunity to poliovirus is defined as: • For children (< 18 years of age): documentation of receipt of > 4 doses of polio vaccine with a minimum interval of 4 weeks between doses; only 3 doses are needed when the third dose is given on or after the fourth birthday. • For adults (> 18 years of age): documentation of receipt of > 3 doses of polio vaccine with a minimum interval of 4 weeks between doses with documentation of > 1 booster dose. Remember, an individual who has received a primary series consisting of > 3 doses of vaccine AND has received > 1 booster dose does NOT need to receive another dose. Note: • Vaccinating an exposed individual who may be incubating poliovirus is not harmful. • Immune globulin (IG) has been found to be of no value as postexposure prophylaxis and is not recommended. (If the use of OPV for a mass vaccination campaign to control a polio outbreak in the U.S. is indicated, the CDC will advise the IDPH on how to obtain an emergency supply of OPV, who should receive OPV, and any other pertinent control measures.) Apply precautions and isolate/exclude as follows:

Page 5

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04 Poliomyelitis
5 • Case: Place on enteric precautions and exclude for 6 weeks after onset or until virus can no longer be recovered from feces (the number of negative specimens needed will be determined on a case-by-case basis). • Contacts: Administer IPV; do not exclude. Surveillance Active surveillance for acute flaccid paralysis and other symptoms of polio infection should continue for at least 2 incubation periods (i.e., up to 70 days) beyond the onset of the last case in an area. D. Preventive Measures Vaccination, including routine childhood vaccination, catch-up vaccination of adolescents, and targeted vaccination of high-risk adult groups, is the best preventive measure against polio. Good personal hygiene (particularly proper handwashing) is also very important.

Routine Polio Childhood Immunization Recommendations An all-IPV polio immunization schedule is now the recommended schedule. OPV is no longer recommended and is not available in the U.S. Four doses of IPV are usually needed to complete the primary series: doses are recommended at ages 2 months, 4 months, 6-18 months, and 4-6 years. At least 28 days are needed between doses, although a 6-8 week interval is preferred between doses 2 and 3 and a 6-month interval is preferred between doses 3 and 4. Only 3 doses are needed when the third dose is given on or after the fourth birthday. Polio vaccine is not routinely recommended for those > 18 years unless there is potential for exposure. Polio Vaccine and Adults Routine vaccination of persons >18 years of age residing in the U.S. is not necessary. However, polio vaccination is indicated for the following groups: • Laboratory workers who handle poliovirus; • Healthcare workers caring for polio patients; • Persons traveling to regions of the world where polio is endemic or epidemic. Polio Vaccination and Travel In assessing the risk to a traveler for polio transmission, healthcare providers are urged to determine first if their patients will truly be traveling to a polio endemic or epidemic area, including Afghanistan, Egypt, India, Niger, Nigeria, and Pakistan. Go <<http://www.cdc.gov/travel>> to obtain information on the risk of transmission of poliovirus in specific countries. Or contact the CDC's Traveler's Health Office at (877) 394-8747. If travel to a polio-endemic or epidemic region is anticipated, please review the patient's history of polio immunization. Ninety percent or more of vaccine recipients develop protective immunity to all three poliovirus types after two doses, and at least 99% are immune following three doses. • If the patient has received a complete primary series of > 3 doses of polio vaccine, administer a booster dose of IPV. Remember, a single booster dose is all that is needed. • If the patient is unimmunized or partially immunized, follow an accelerated schedule to complete as much of the series as possible before departure, as outlined in the table below: Weeks Available Accelerated IPV Schedule* > 8 weeks 3 doses, given 4 weeks apart 4-7 weeks 2 doses, given 4 weeks apart < 4 weeks 1 dose *1stdose may be given as early as 6 weeks of age

Page 6

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04 Poliomyelitis 64) ADDITIONAL INFORMATION The following is the formal CSTE surveillance case definition for polio. It is provided for your information only. It is not necessary to use this information for reporting or investigating a case. (CDC case definitions are used by the state health department and CDC to maintain uniform standards for national reporting.) For reporting to the state, always use the criteria outlined in Section 2) A. Clinical case definition for paralytic poliomyelitis: Acute onset of a flaccid paralysis of one or more limbs with decreased or absent tendon reflexes in the affected limbs, without other apparent cause, and without sensory or cognitive loss. Case classification Probable: A case that meets the clinical case definition. Confirmed: A case that meets the clinical case definition and in which the patient has a neurologic deficit 60 days after onset of initial symptoms, has died, or has unknown follow-up status. Laboratory criteria for diagnosis The following is the formal CDC surveillance case definition for polio. It is provided for your information only. It is not necessary to use this information for reporting or investigating a case. (CDC case definitions are used by the state health department and CDC to maintain uniform standards for national reporting.)

Clinical case definition for paralytic poliomyelitis: Acute onset of a flaccid paralysis of one or more limbs with decreased or absent tendon reflexes in the affected limbs, without other apparent cause, and without sensory or cognitive loss. Case classification Probable: A case that meets the clinical case definition. Confirmed: A case that meets the clinical case definition and in which the patient has a neurologic deficit 60 days after onset of initial symptoms, has died, or has unknown follow-up status. The likelihood of poliovirus isolation is highest from stool specimens, intermediate from pharyngeal swabs, and very low from blood or spinal fluid. The isolation of poliovirus from stool specimens contributes to the diagnostic evaluation but does not constitute proof of a causal association of such viruses with paralytic poliomyelitis. Isolation of virus from the cerebrospinal fluid (CSF) is diagnostic but is rarely accomplished. To increase the probability of poliovirus isolation, at least two stool specimens and two throat swabs should be obtained 24 hours apart from patients with suspected poliomyelitis as early in the course of the disease as possible (i.e., immediately after poliomyelitis is considered as a possible differential diagnosis), but ideally within the first 14 days after onset of paralytic disease. Specimens should be sent to the University Hygienic Lab.

Page 7

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04Poliomyelitis
7References American Academy of Pediatrics. Red Book 2003: Report of the Committee on Infectious Diseases, 26th Edition. Illinois, Academy of Pediatrics, 2003. CDC. Case Definitions for Infectious Conditions Under Public Health Surveillance. MMWR. 1997; 46:RR-10. CDC. Epidemiology & Prevention of Vaccine-Preventable Diseases: The Pink Book, 8th Edition. CDC, January 2004. CDC. Vaccine-Preventable Disease Surveillance Manual, 3rd Edition, 2002. CDC, 1999. Chin, J., ed. Control of Communicable Diseases Manual, 17th Edition. Washington, DC, American Public Health Association, 2000. IDPH. Public Health (641) Chapter 1, Notification and Surveillance of Reportable Communicable and Infectious Diseases, Poisonings and Conditions (Printed April 2004). Additional Resources “Manual for the Surveillance of VPDs”, Chapter 10: Poliomyelitis. Available at the following web site: http://www.cdc.gov/nip/publications/surv-manual/chpt10_polio.pdf.

Page 8

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04Poliomyelitis
8

Page 9

Iowa Department of Public Health Iowa Dept. of Public Health Rev. 8/04Poliomyelitis
9FACT SHEETPOLIOMYELITISInformation for Health Professionals (Polio) What is poliomyelitis? A viral infection most often recognized by the acute onset of flaccid paralysis. Poliovirus infection occurs in the GI track with spread to the regional nodes and in a minority of cases, to the nervous system. Who gets poliomyelitis? Prior to the widespread use of polio vaccine, poliomyelitis occurred worldwide. Polio was epidemic in the U.S. for the first half of the 20th century with over 20,000 cases of paralytic disease in 1952. The first inactivated poliovirus vaccine (IPV) was introduced in 1955,

monovalent oral poliovirus vaccine (OPV) in 1961, trivalent in 1963, and enhanced inactivated poliovirus vaccine (eIPV) in 1987. After the introduction of vaccination, the reported number of cases of poliomyelitis in the U.S. dropped to <100 in 1965 and <10 cases in 1973. The last cases of indigenously-transmitted wild-type poliovirus in the U.S. were in 1979. The last case of wild-type polio disease in the Western Hemisphere was detected in Peru in 1991. The Western Hemisphere was declared free from indigenous wild-type poliovirus transmission in 1994. Almost all of the world is now considered polio-free. Worldwide efforts to eradicate polio in countries where the disease is still endemic are underway. Strategies include: (1) achieving and maintaining high vaccination coverage among infants < 1 year old; (2) developing sensitive surveillance systems for AFP and a laboratory network; (3) conducting National Immunization Days; (4) and conducting “mopping-up” campaigns to directly target geographic areas known to be high risk for polio transmission. The number of countries where poliovirus continues to be isolated has decreased substantially, with Afghanistan, Egypt, India, Niger, Nigeria, and Pakistan remaining the major areas of wild-type virus circulation. How is the poliomyelitis spread? The principal mode of transmission is person-to-person by the fecal-oral or oral-oral route, with the fecal-oral route predominating. Transmission via oral secretions, such as saliva, is possible and may account for some cases. In rare instances, the virus may be transmitted by contaminated sewage or water. Asymptomatic individuals, especially children, comprise a significant source of infections. No reliable evidence of spread by insects exists. No long-term carrier state is known. In temperate climates, poliovirus infections are most common in the summer and fall. What are the symptoms? Infection with poliovirus results in a spectrum of manifestations. The overwhelming majority of infections (95%) are clinically inapparent. Some 4 – 8% of infected individuals will experience non-specific viral symptoms, such as a low-grade fever, headache, sore throat, nausea, abdominal pain, constipation, diarrhea, and/or vomiting (abortive disease). Some 1–5% of infections will result in aseptic meningitis, involving stiffness of the back, neck and/or legs, at times with paresthesias, a few days after the minor illness has resolved. Only about 0.1–1% of infections will progress to acute flaccid paralysis (AFP) with loss of reflexes in the involved limbs, usually with fever present (paralytic poliomyelitis). Please note, today in the U.S., the most common cause of AFP is Guillain-Barré syndrome. How soon do the symptoms appear? Progression to paralytic poliomyelitis usually occurs within 2 – 4 days and rarely continues after the fever subsides. How long can an infected person spread the virus? The period of communicability is not precisely defined. It appears greatest 7-10 days before and after onset of clinical symptoms, when poliovirus is present in the throat and excreted in the highest quantities in the feces. Poliovirus can continue to be shed in the feces for 4 to 6 weeks. Rarely, excretion of poliovirus >6 months after infection has been found in asymptomatic, immunodeficient persons. Poliovirus can be found in throat secretions as early as 36 hours and in the feces 72 hours after exposure to infection in both symptomatic and asymptomatic cases.

Also, attempt to identify the route of introduction of poliovirus into the community. To identify these groups, think in terms of “zones of exposure” and consider members of the following groups: • Household members • School/child care associates (students/attendees and staff) • Staff and patients at medical facility where patient was cared for, especially if there was the potential for direct contact with feces or oral secretions • Religious/social groups • Sports teams and other extracurricular groups • Bus mates • Close friends • Travelers from polio-endemic regions such as Afghanistan, Egypt, India, Niger, Nigeria, and Pakistan • Any other persons who may have come in direct contact with the case’s feces or oral secretions

What are poliomyelitis isolation guidelines? Place case on enteric precautions for six weeks after onset of symptoms or until poliovirus can no longer be recovered from feces (the number of negative specimens needed will be determined by the IDPH on a case-by-case basis). Can a person get poliomyelitis again? Adults who contract paralytic poliomyelitis during childhood may develop the postpolio syndrome 30 to 40 years later. Postpolio syndrome is characterized by slow onset of muscle pain and exacerbation of weakness. What is the treatment for poliomyelitis? None. Supportive treatment and attention during the acute illness to the complications of paralysis requires expert knowledge and equipment, especially for patients in need of respiratory assistance. Is there a vaccine to prevent poliomyelitis? Yes, there is a vaccine to protect against poliomyelitis. An all-IPV polio immunization schedule is now the recommended schedule. OPV is no longer recommended and is not available in the U.S. Four doses of IPV are usually needed to complete the primary series: doses are recommended at ages 2 months, 4 months, 6-18 months, and 4-6 years. At least 28 days are needed between doses, although a 6-8 week interval is preferred between doses 2 and 3 and a 6-month interval is preferred between doses 3 and 4. Only 3 doses are needed when the third dose is given on or after the fourth birthday. Polio vaccine is not routinely recommended for those > 18 years unless there is potential for exposure. Routine vaccination of persons >18 years old residing in the U.S. is not necessary. However, polio vaccination is indicated for the following groups: • Laboratory workers who handle poliovirus; • Healthcare workers caring for polio patients; • Persons traveling to regions of the world where polio is endemic or epidemic.